## 2020/2021 Standard Wheat Test

(13 Varieties x 8 Locations)



MS	Variety	AvgYld	MOIST	TWT	Carroll	Fayette	Gibson	Henry	Madison	Moore	Weakley	WTREC	
		bu/ac	%	lb/bu	20-Nov	9-Nov	14-Nov	4-Nov	13-Nov	4-Nov	5-Nov	14-Oct	% ≥ Avg.
А	USG 3329	100.8	14.2	58.1	70	124	67	89	132	108	92	124	88
AB	Progeny #BUSTER	98.5	14.3	59.4	70	105	58	93	132	101	104	125	88
AB	Dyna-Gro 9151	98.4	13.8	60.4	68	104	60	97	136	116	81	126	75
ABC	Progeny #CHAD	97.1	14.5	57.5	82	88	71	99	125	100	89	123	75
ABCD	Progeny #BLAZE	96.9	13.7	58.9	61	109	63	94	133	106	86	123	75
ABCD	Dyna-Gro 9002	95.3	13.9	59.4	62	112	58	87	130	105	87	122	50
ABCDE	Dyna-Gro 9172	94.1	13.6	60.1	58	99	72	83	130	100	88	123	63
BCDE	Dyna-Gro 9692	92.7	13.7	59.3	70	79	62	91	135	94	91	118	63
BCDE	AgriPro SY Viper	91.5	13.9	59.1	61	77	55	100	131	102	78	128	38
BCDE	AgriPro SR 1421	91.1	14.3	56.5	66	92	52	96	134	89	87	113	38
CDE	USG 3472	90.2	13.5	58.4	53	97	51	87	133	90	95	117	38
DE	USG 3562	89.0	14.4	59.0	66	90	68	82	124	104	69	109	38
Е	AgriPro SY Richie	86.8	13.8	58.3	65	68	67	82	131	78	85	117	25
	Averages	94.0	14.0	58.8	65.6	95.6	61.7	90.8	131.2	99.5	87.1	120.7	

Yields have been adjusted to 13.5% moisture. Each variety was evaluated in a large strip-plot at each

location, thus each county test was considered as one replication of the test in calculating the average yield

Statistical analysis conducted at 95% confidence interval. Varieties with the same MS letter are not significantly different.

Official test weight of No. 2 wheat=58 lbs/bu. TWT = Avg. Test Wt. lbs./bu @ 7 locations.

 $\% \ge$  Avg. is the % of locations that variety is at or above the field average.

Programs in agriculture and natural resources, 4-H youth development, family and consumer sciences, and resource development

University of Tennessee Institute of Agriculture, U.S. Department of Agriculture and county governments cooperating.

UT Extension provides equal opportunities in programs and employment.