

Table A-3 cont.

Hybrid [†]	Herbicide Pkg. [‡]	Insect Pkg. [‡]	Test Weight (lbs/bu)			Protein (%)			Oil (%)			Starch (%)		
			1 yr	2 yr	3 yr	1 yr	2 yr	3 yr	1 yr	2 yr	3 yr	1 yr	2 yr	3 yr
Dekalb DKC68-35	RR	VT2P	59.1 AB			8.6 BC			3.5 E			73.1 A		
Revere 1839	RR	TRE	55.4 E			8.4 C			3.8 B-D			72.4 AB		
LG 67C07	RR	VT2P	58.6 B	56.9 A		8.8 A-C	8.3 A		3.7 C-E	3.7 A		72.7 A	72.6 A	
LG 68C18	RR	VT2P	57.7 CD			9.2 A			4.1 A			70.8 C		
Dyna-Gro D57VC53	RR	VT2P	59.6 A	57.4 A		9.1 AB	8.5 A		4.0 AB	3.9 A		72.5 AB	72.4 A	
Progeny 9117	RR	VT2P	57.0 D	54.0 B	55.0 B	8.7 A-C	8.5 A	8.6 A	3.8 B-D	3.8 A	3.9 A	71.3 BC	72.0 A	71.9 A
Dekalb DKC70-45	RR	VT2P	58.4 BC			9.0 AB			3.7 DE			72.2 AB		
Progeny 2118	RR	VT2P	59.6 A	55.9 AB	57.3 A	8.3 C	8.3 A	8.6 A	3.9 A-C	3.7 A	3.8 A	72.4 AB	72.5 A	72.5 A
Average			58.2	56.0	56.2	8.7	8.4	8.6	3.8	3.8	3.8	72.2	72.4	72.2
Standard Error			0.3	2.8	2.4	0.2	0.4	0.3	0.1	0.1	0.1	0.4	0.3	0.3
L.S.D. ^{.05}			0.8	2.3	2	0.6	N.S.	N.S.	0.2	N.S.	N.S.	1.2	N.S.	N.S.
C.V.			0.8	3.4	2.7	3.8	5.2	4.9	3.1	4.0	3.8	0.9	1.2	0.8

[†] Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

[‡] For a full description of abbreviated biotech traits, see table 18.

[§] All yields are adjusted to 15.5% moisture.

^{||} Lodging values do not typically follow a normal distribution, therefore statistical tests to compute LSD were not performed and only mean values are reported.

Values highlighted in light orange are above average for a given trait, MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.