

**Table 19. Yields of 18 Maturity Group IV Late (4.6-4.9) Glyphosate / Dicamba tolerant soybean varieties in 12 County Standard Tests in Tennessee during 2023<sup>†</sup>.**

MS† Avg. Yield	Variety	Avg. Yield <sup>§</sup> (bu/acre)	Avg. Moisture (%)	Croc	Dyer	Frank	Gibs	Giles	Hard	Hardin	Henr	Laud	Mad	Perr	Weak
				6/27	6/1	5/31	5/10	5/30	5/3	6/1	6/26	5/27	6/24	5/24	5/10
A	Pioneer P47A64X	<b>71.8</b>	10.7	<b>80</b>	56	81	<b>88</b>	59	75	59	63	45	<b>98</b>	98	56
AB	NK 46-B4XFS	69.8	10.7	73	66'	77	82	49	<b>87</b>	<b>66</b>	63	39	94	97	44
ABCD	Revere 4795XS*	68.8	10.2	74'	65	80	84	<b>68</b>	64	56	62	41	83	96	51
ABC	Revere 4826XFS	68.8	10.6	74	65	82	80	67	80	60	51	37	88	89	53
ABCD	Dyna-Gro S49XF43S	67.6	11.0	76	63'	81	80	47	74	57	62	33	94	99	46
ABCD	USG 7474XFS	66.4	10.8	70	57	77	78	43	79	53	63	38	88	86	<b>65</b>
BCD	Revere 4806XS	66.2	<b>11.1</b>	68	69	77	75	67	67	63	58	33	85	89	43
BCD	Asgrow 48XF3	66.2	10.4	76	63'	74	75	37	70	60	60	42	82	91	65
BCD	NK 48-A8XF	65.9	10.8	71	<b>73</b>	80	78	40	67	56	63	39	88	90	43
BCD	Armor 46-F76	65.6	10.6	67	68	69	79	43	74	50	<b>64</b>	43	81	90	59
BCD	Pioneer P46A90LX	65.5	10.6	73	57	79	82	43	57	51	62	<b>53</b>	91	91	47
BCD	Asgrow 49XF3	65.3	10.8	75	68	79	71	48	69	56	61	24	92	81	60
BCD	Progeny 4691XFS	64.5	10.8	68	68	68	75	42	75	64	61	34	88	93	37
BCD	USG 7463XF	64.4	11.0	69	58	75	77	66	69	56	57	45	81	90	32
CD	Armor 46-F96	64.0	10.1	67	54	77	80	44	73	50	61	38	88	89	47
D	Asgrow 47XF2	63.4	10.5	63	64	82	81	40	48	55	53	37	80	<b>103</b>	55
D	USG 7461XFS	63.3	10.7	70	51	<b>83</b>	80	35	73	63	60	37	83	89	36
D	Dyna-Gro S47XF23S	63.1	10.4	62	63	73	79	46	72	45	61	35	85	86	51
<b>Average</b>		<b>66.1</b>	<b>10.6</b>	<b>71</b>	<b>62</b>	<b>78</b>	<b>79</b>	<b>49</b>	<b>71</b>	<b>57</b>	<b>60</b>	<b>38</b>	<b>87</b>	<b>91</b>	<b>49</b>

Data Provided by Ryan Blair, Ext. Area Specialist, Grain and Cotton Variety Testing, and Extension agents in counties shown above.

† Varieties that have any MS letter in common are not significantly different at the 5% level of probability. Values highlighted in light orange are above average for a given trait, MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.

\* Asterisks after a name indicate the number of preceding consecutive years in the top-performing "A" group.

§ All yields are adjusted to 13% moisture.

County Locations include: Crockett, Dyer, Franklin, Gibson, Giles, Hardeman, Hardin, Henry, Lauderdale, Madison, Perry, Weakley