

**Table A-16. Mean yield and agronomic traits of 10 early-season (<114 DAP) corn hybrids evaluated in small plot replicated trials with irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2024.**

Hybrid <sup>†</sup>	Herbicide Pkg <sup>‡</sup>	Insect Pkg. <sup>‡</sup>	Avg. Yield <sup>§</sup> (bu/ac)	Moisture at Harvest (%)
Pioneer P13777PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	259 A	15.8 C
Innvictis A1292 VT2P	RR	VT2P	252 AB	15.9 BC
Dekalb DKC 111-35 VT2P RIB	RR	VT2P	252 AB	15.7 CD
Revere 113-T4C	RR	CB, VP	246 A-C	15.8 C
Dyna-Gro D53VC54 RIB	RR	VT2P	239 BC	17.1 AB
Progeny PGY 2010 TRE	RR	TRE	238 BC	14.4 E
Dyna-Gro D51VC95 RIB	RR	VT2P	236 B-D	15.2 C-E
Innvictis A1072 VT2P RIB	RR	VT2P	234 CD	14.6 DE
Pioneer P13841PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	232 CD	14.9 C-E
Great Heart Seed HT-7360 VT2	RR	VT2P	220 D	17.2 A
<b>Trial Average</b>			241	15.7
<b>Trial Standard Error</b>			7	0.4
<b>Trial L.S.D.<sub>.05</sub></b>			17	1.2
<b>Trial C.V.</b>			4	4

† Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

\* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

‡ For a full description of abbreviated biotech traits, see table 18.

§ All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.