

Table A-18. Mean yield and agronomic traits of nine full-season (>116 DAP) corn hybrids evaluated in small plot replicated trials with irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2024.

Hybrid [†]	Herbicide Pkg [‡]	Insect Pkg. [‡]	Avg. Yield [§] (bu/ac)	Moisture at Harvest (%)
Dekalb DKC 68-35 VT2P*	RR	VT2P	254 A	16.6 A
Revere 1839 TC*	RR	TRE	247 AB	15.1 D
Integra 6915 TRE	RR	TRE	239 A-C	15.2 CD
Innictis A1993 T	RR	TRE	233 B-D	15.1 D
Innictis A1792 T	RR	TRE	230 B-D	16.2 AB
Dyna-Gro D58VC74 RIB	RR	VT2P	224 C-E	15.8 BC
Progeny PGY 2118 VT2P	RR	VT2P	220 DE	16.5 A
Progeny PGY 9117 VT2P	RR	VT2P	219 DE	15.6 B-D
Pioneer P17677YHR	RR, LL	YGCB, HX1	209 E	15.8 BC
Trial Average			231	15.8
Trial Standard Error			6	0.2
Trial L.S.D._{.05}			18	0.6
Trial C.V.			5	2

[†] Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

[‡] For a full description of abbreviated biotech traits, see table 18.

[§] All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.