

Table A-19. Mean yield and agronomic traits of 10 early-season (<114 DAP) corn hybrids evaluated in small plot replicated trials without irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2024.

Hybrid [†]	Herbicide Pkg [‡]	Insect Pkg. [‡]	Avg. Yield [§] (bu/ac)	Moisture at Harvest (%)
Dekalb DKC 111-35 VT2P RIB	RR	VT2P	241 A	14.6 AB
Innvictis A1292 VT2P	RR	VT2P	239 A	15.1 A
Pioneer P13777PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	231 AB	14.8 AB
Pioneer P13841PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	228 AB	13.3 CD
Revere 113-T4C	RR	CB, VP	227 AB	14.0 BC
Progeny PGY 2010 TRE	RR	TRE	217 BC	12.7 D
Dyna-Gro D53VC54 RIB	RR	VT2P	215 BC	15.3 A
Dyna-Gro D51VC95 RIB	RR	VT2P	213 BC	13.2 CD
Innvictis A1072 VT2P RIB	RR	VT2P	212 BC	12.7 D
Great Heart Seed HT-7360 VT2	RR	VT2P	207 C	15.4 A
Trial Average			223	14.1
Trial Standard Error			8	0.3
Trial L.S.D._{.05}			20	0.9
Trial C.V.			5	4

† Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

‡ For a full description of abbreviated biotech traits, see table 18.

§ All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highlighted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.