

Table A-21. Mean yield and agronomic traits of nine full-season (>116 DAP) corn hybrids evaluated in small plot replicated trials without irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2024.

Hybrid [†]	Herbicide Pkg [‡]	Insect Pkg. [‡]	Avg. Yield [§] (bu/ac)	Moisture at Harvest (%)
Revere 1839 TC*	RR	TRE	233 A	14.0 E
Dekalb DKC 68-35 VT2P*	RR	VT2P	230 A	15.0 B-D
Integra 6915 TRE	RR	TRE	228 A	14.2 DE
Innictis A1993 T	RR	TRE	227 A	14.3 C-E
Innictis A1792 T	RR	TRE	224 AB	14.9 B-E
Dyna-Gro D58VC74 RIB	RR	VT2P	222 AB	15.2 BC
Progeny PGY 9117 VT2P	RR	VT2P	220 A-C	15.3 AB
Progeny PGY 2118 VT2P	RR	VT2P	209 BC	16.2 A
Pioneer P17677YHR	RR, LL	YGCB, HX1	203 C	15.0 B-D
Trial Average			222	14.9
Trial Standard Error			7	0.3
Trial L.S.D._{.05}			18	0.9
Trial C.V.			5	4

[†] Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

[‡] For a full description of abbreviated biotech traits, see table 18.

[§] All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.