

**Table A-25. Mean yield and agronomic traits of 10 early-season (<114 DAP) corn hybrids evaluated in small plot replicated trials with irrigation at AgriCenter International in Memphis, Tennessee during 2024.**

Hybrid <sup>†</sup>	Herbicide Pkg <sup>‡</sup>	Insect Pkg. <sup>‡</sup>	Avg. Yield <sup>§</sup> (bu/ac)	Moisture at Harvest (%)	Test Weight (lbs/bu)
Great Heart Seed HT-7360 VT2	RR	VT2P	204 A	14.0 A	55 A
Dekalb DKC 111-35 VT2P RIB	RR	VT2P	185 A	14.3 A	53 A
Innvictis A1292 VT2P	RR	VT2P	181 A	14.3 A	54 A
Revere 113-T4C	RR	CB, VP	179 A	14.7 A	53 A
Dyna-Gro D51VC95 RIB	RR	VT2P	178 A	13.6 A	52 A
Innvictis A1072 VT2P RIB	RR	VT2P	155 A	13.7 A	57 A
Dyna-Gro D53VC54 RIB	RR	VT2P	152 A	13.7 A	56 A
Progeny PGY 2010 TRE	RR	TRE	150 A	14.0 A	57 A
Pioneer P13841PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	147 A	13.2 A	53 A
Pioneer P13777PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	145 A	14.7 A	56 A
<b>Trial Average</b>			168	14.0	54
<b>Trial Standard Error</b>			15	0.6	2
<b>Trial L.S.D.<sub>.05</sub></b>			N.S.	N.S.	N.S.
<b>Trial C.V.</b>			16	7	5

† Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

\* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

‡ For a full description of abbreviated biotech traits, see table 18.

§ All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.