

Table 19. Yields of 8 full-season (>116 DAP) Roundup / stacked corn hybrids in 13 County Standard Tests in Tennessee during 2025.‡

MS† Avg. Yield	Hybrid*	Avg. Yield§ bu/acre	Avg. Moisture %	Avg. Test Weight lbs/bu	Henry Henry												
					Chest 4/17	Crock 5/9	Gibs 4/18	Giles 4/25	Harde 5/1	Hayw 4/15	B 4/15	T 4/24	Loud 4/17	Mad 6/3	Mont 5/6	Obion 4/18	Weak 4/30
A	DynaGro 60TC45*	227.8	15.1	59.7	240	280	193	267	223	252	297	247	144	107	150	300	262
A	Beck's 6973 TCV2P	225.6	15.4	59.7	236	275	188	275	246	253	277	239	126	120	154	289	255
A	Dekalb 68-35 VT2P*	221.8	14.4	60.8	233	253	198	271	232	244	277	226	156	95	144	292	263
AB	Gateway 3919 TRE	218.8	15.4	58.3	243	250	190	263	231	238	283	238	124	105	111	302	265
BC	Agrigold 647-79 VT2P	212.3	14.6	60.8	243	242	186	260	228	236	264	218	143	113	132	262	230
BC	Pioneer P18216 PWE	211.7	14.9	59.5	218	247	172	268	209	238	273	217	144	106	141	280	237
C	DynaGro 58VC74	203.7	14.7	60.5	203	245	180	252	210	201	245	222	112	109	169	260	245
C	Pioneer P17677 YHR	203.3	14.1	61.0	198	234	178	247	198	227	251	205	120	118	153	280	235
Average		216	15	60	227	253	186	263	222	236	271	227	134	109	144	283	249

‡ Data Provided by Ryan Blair, Ext. Area Specialist, Grain and Cotton Variety Testing, and Extension agents in counties shown above.

† Hybrids that have any MS letter in common are not significantly different in yield at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

§ All yields are adjusted to 15.5% moisture.

Highlighted cells indicate hybrids that were above average and bold/underline values indicate the top yield, within a location.

County locations include: Chester, Crockett, Gibson, Giles, Hardeman, Haywood, Henry (2 locs), Loudon, Madison, Montgomery, Obion, & Weakley.