

Table A-15. Mean yield and agronomic traits of 11 full-season (>116 DAP) corn hybrids evaluated in small plot replicated trials without irrigation at the Middle Tennessee AgResearch and Education Center in Spring Hill, Tennessee during 2025.

Hybrid [†]	Herbicide Pkg [‡]	Insect Pkg. [‡]	Avg. Yield [§] (bu/ac)	Moisture at Harvest (%)	Plant Height (in.)	Ear Height (in.)	Lodging [¶] (%)
Dekalb DKC 119-30	RR	VT4P	216 A	15.7 AB	114 E	43 C	0.0
Crow's 5859 TRE	RR	TRE	211 A	14.6 C	122 AB	52 A	0.0
Dekalb DKC 68-35**	RR	VT2P	209 A	15.6 AB	116 DE	43 C	0.0
Innvictis A1792T	RR	TRE	200 A	15.9 A	120 A-D	50 AB	0.0
Revere 1839 TC**	RR	TRE	196 A	15.0 BC	123 AB	53 A	0.0
Integra 6915 TRE*	RR	TRE	191 A	14.5 C	119 B-D	48 AB	0.0
Innvictis A1993T*	RR	TRE	191 A	15.1 A-C	122 A-C	51 A	0.0
Dyna-Gro D60TC45RIB	RR	TRE	186 A	14.7 C	121 A-C	51 A	0.0
Pioneer P17677YHR	RR, LL	YGCB, HX1	183 A	14.3 C	124 A	49 AB	0.0
Progeny PGY 2419TRE	RR	TRE	177 A	14.7 C	119 B-D	51 A	0.0
Pioneer P18216PWE	RR, LL, ENL, FOP	VT2P, HX1	167 A	14.3 C	118 C-E	46 BC	0.0
Trial Average			193	14.9	120	49	0.0
Trial Standard Error			17	0.4	2	2	0.0
Trial L.S.D._{.05}			N.S.	0.9	4	4	.
Trial C.V.			14	3	2	5	0.0

[†] Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

[‡] For a full description of abbreviated biotech traits, see table 4.

[§] All yields are adjusted to 15.5% moisture.

[¶] Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.