

**Table A-16. Mean yield and agronomic traits of 14 early-season (<114 DAP) corn hybrids evaluated in small plot replicated trials with irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2025.**

Hybrid <sup>†</sup>	Herbicide Pkg <sup>‡</sup>	Insect Pkg. <sup>‡</sup>	Avg. Yield <sup>§</sup> (bu/ac)	Moisture at Harvest (%)	Test Weight (lbs/bu)
Pioneer P13777PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	261 A	12.7 AB	60 B-D
Dekalb DKC 111-35 RIB*	RR	VT2P	257 A	12.1 B-E	61 A
Great Heart 7335TRC	RR	TRE	257 A	13.1 A	61 AB
Augusta A4862	RR, LL	BT	256 A	12.2 B-E	58 G
Dyna-Gro D52TC66RIB	RR	TRE	252 A	12.4 A-C	60 C-E
Innvictis A1292VT2P	RR	VT2P	246 A	11.9 C-E	61 A
Pioneer P11616PWE	RR, LL, ENL, FOP	VT2P, HX1	240 AB	11.5 E	60 BC
Innvictis A1254T	RR	TRE	238 AB	12.4 A-D	59 D-F
Great Heart 7210TRC	RR	TRE	234 AB	12.8 AB	59 C-E
Innvictis A1312VT2P*	RR	VT2P	220 BC	13.0 A	58 FG
Cane Run Enterprises CRE-F12	none	none	218 BC	11.5 E	59 EF
Cane Run Enterprises CRE-Z99	none	none	207 CD	11.6 DE	59 EF
Cane Run Enterprises CRE-135B	none	none	189 D	11.5 E	58 FG
Cane Run Enterprises CRE-48N	none	none	183 D	12.4 A-C	58 G
<b>Trial Average</b>			233	12.2	59
<b>Trial Standard Error</b>			11	0.3	0
<b>Trial L.S.D.<sub>.05</sub></b>			24	0.7	1
<b>Trial C.V.</b>			6	3	1

<sup>†</sup> Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

\* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

<sup>‡</sup> For a full description of abbreviated biotech traits, see table 4.

<sup>§</sup> All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.