

Table A-17. Mean yield and agronomic traits of 15 medium-season (114-116 DAP) corn hybrids evaluated in small plot replicated trials with irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2025.

Hybrid [†]	Herbicide Pkg [‡]	Insect Pkg. [‡]	Avg. Yield [§] (bu/ac)	Moisture at Harvest (%)	Test Weight (lbs/bu)
Crow's 5444 VT2P	RR	VT2P	251 A	12.2 C-E	61 BC
Innvictis A1542T*	RR	TRE	249 A	12.2 C-E	60 D-F
Innvictis A1414T	RR	TRE	239 AB	12.4 C-E	61 CD
Revere 1627 TC***	RR	TRE	238 AB	12.4 C-E	61 BC
Pioneer P14364PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	238 AB	12.3 C-E	59 G
Dekalb DKC 64-22*	RR	VT2P	236 A-C	11.9 E	63 A
Progeny PGY 2314TRE**	RR	TRE	236 A-C	13.0 BC	61 CD
Integra 6624 TRE	RR	TRE	234 A-C	12.1 C-E	60 EF
Revere 114-P35*	RR	CB	232 A-C	13.6 AB	60 FG
Innvictis A1551VT2P	RR	VT2P	230 A-C	11.9 E	60 EF
Dyna-Gro D55TC86RIB	RR	TRE	229 A-C	12.0 DE	61 C-E
Great Heart 7451VT2	RR	VT2P	225 BC	12.9 B-D	63 A
Dekalb DKC 114-99	RR	VT4P	222 B-D	12.5 C-E	62 B
Innvictis X1485PWE	RR	PWE	215 CD	12.1 DE	61 C-F
Cane Run Enterprises CRE-2020	none	none	200 D	14.2 A	58 H
Trial Average			232	12.5	61
Trial Standard Error			10	0.3	0
Trial L.S.D._{.05}			22	0.8	1
Trial C.V.			6	4	1

[†] Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

[‡] For a full description of abbreviated biotech traits, see table 4.

[§] All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highlighted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.