

Table A-18. Mean yield and agronomic traits of 11 full-season (>116 DAP) corn hybrids evaluated in small plot replicated trials with irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2025.

Hybrid [†]	Herbicide Pkg [‡]	Insect Pkg. [‡]	Avg. Yield [§] (bu/ac)	Moisture at Harvest (%)	Test Weight (lbs/bu)
Integra 6915 TRE*	RR	TRE	232 A	11.7 C-E	60 CD
Dekalb DKC 119-30	RR	VT4P	225 AB	13.3 A	61 B
Crow's 5859 TRE	RR	TRE	225 A-C	11.4 E	60 D
Pioneer P18216PWE	RR, LL, ENL, FOP	VT2P, HX1	224 A-C	12.5 B	60 D
Innvictis A1993T*	RR	TRE	219 B-D	11.7 C-E	60 D
Dekalb DKC 68-35**	RR	VT2P	218 B-D	12.2 BC	61 BC
Pioneer P17677YHR	RR, LL	YGCB, HX1	217 B-D	11.7 DE	61 B
Progeny PGY 2419TRE	RR	TRE	215 B-D	11.8 C-E	60 D
Dyna-Gro D60TC45RIB	RR	TRE	213 CD	11.8 C-E	60 D
Revere 1839 TC**	RR	TRE	211 D	12.0 B-D	60 D
Innvictis A1792T	RR	TRE	209 D	12.2 B-D	63 A
Trial Average			219	12.0	61
Trial Standard Error			7	0.2	0
Trial L.S.D._{.05}			12	0.6	0
Trial C.V.			3	3	0

† Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

‡ For a full description of abbreviated biotech traits, see table 4.

§ All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.