

Table A-19. Mean yield and agronomic traits of 14 early-season (<114 DAP) corn hybrids evaluated in small plot replicated trials without irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2025.

Hybrid [†]	Herbicide Pkg [‡]	Insect Pkg. [‡]	Avg. Yield [§] (bu/ac)	Moisture at Harvest (%)	Test Weight (lbs/bu)
Pioneer P13777PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	253 A	12.1 A	59 A-C
Cane Run Enterprises CRE-135B	none	none	251 A	11.8 A	60 AB
Cane Run Enterprises CRE-48N	none	none	249 A	12.2 A	59 A-C
Innvictis A1292VT2P	RR	VT2P	247 A	12.1 A	60 A-C
Great Heart 7335TRC	RR	TRE	245 A	12.7 A	60 AB
Innvictis A1254T	RR	TRE	244 A	12.4 A	59 CD
Great Heart 7210TRC	RR	TRE	243 A	12.0 A	60 A
Cane Run Enterprises CRE-F12	none	none	243 A	12.3 A	60 A
Dyna-Gro D52TC66RIB	RR	TRE	242 A	12.5 A	60 A-C
Augusta A4862	RR, LL	BT	240 A	11.9 A	58 D
Dekalb DKC 111-35 RIB*	RR	VT2P	238 A	11.9 A	60 A-C
Innvictis A1312VT2P*	RR	VT2P	236 A	11.9 A	60 A-C
Pioneer P11616PWE	RR, LL, ENL, FOP	VT2P, HX1	230 A	12.1 A	59 A-D
Cane Run Enterprises CRE-Z99	none	none	225 A	11.5 A	59 B-D
Trial Average			242	12.1	59
Trial Standard Error			9	0.4	1
Trial L.S.D._{.05}			N.S.	N.S.	1
Trial C.V.			6	4	1

[†] Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

[‡] For a full description of abbreviated biotech traits, see table 4.

[§] All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.