

**Table A-20. Mean yield and agronomic traits of 15 medium-season (114-116 DAP) corn hybrids evaluated in small plot replicated trials without irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2025.**

Hybrid <sup>†</sup>	Herbicide Pkg <sup>‡</sup>	Insect Pkg. <sup>‡</sup>	Avg. Yield <sup>§</sup> (bu/ac)	Moisture at Harvest (%)	Test Weight (lbs/bu)
Dekalb DKC 114-99	RR	VT4P	242 A-D	12.6 A	61 AB
Dekalb DKC 64-22*	RR	VT2P	248 A-C	12.2 A	61 AB
Cane Run Enterprises CRE-2020	none	none	214 F	12.8 A	58 EF
Integra 6624 TRE	RR	TRE	240 B-D	12.1 A	60 CD
Pioneer P14364PWUE	RR, LL, ENL, FOP	AVBL, VT2P, HX1	254 A	12.8 A	58 F
Progeny PGY 2314TRE**	RR	TRE	240 B-D	12.6 A	59 DE
Great Heart 7451VT2	RR	VT2P	238 CD	13.0 A	62 A
Innvictis A1414T	RR	TRE	243 A-D	12.7 A	60 CD
Innvictis A1542T*	RR	TRE	253 AB	12.5 A	60 B-D
Innvictis A1551VT2P	RR	VT2P	221 EF	11.8 A	59 DE
Innvictis X1485PWE	RR	PWE	250 A-C	12.1 A	60 B-D
Dyna-Gro D55TC86RIB	RR	TRE	251 A-C	13.0 A	59 CD
Crow's 5444 VT2P	RR	VT2P	245 A-D	11.9 A	60 BC
Revere 114-P35*	RR	CB	255 A	12.7 A	59 DE
Revere 1627 TC***	RR	TRE	233 DE	12.9 A	60 CD
<b>Trial Average</b>			242	12.5	60
<b>Trial Standard Error</b>			6	0.4	0
<b>Trial L.S.D.<sub>.05</sub></b>			12	N.S.	1
<b>Trial C.V.</b>			3	4	1

† Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

\* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

‡ For a full description of abbreviated biotech traits, see table 4.

§ All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highlighted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.