

**Table A-21. Mean yield and agronomic traits of 11 full-season (>116 DAP) corn hybrids evaluated in small plot replicated trials without irrigation at the AgResearch and Education Center at Milan in Milan, Tennessee during 2025.**

Hybrid <sup>†</sup>	Herbicide Pkg <sup>‡</sup>	Insect Pkg. <sup>‡</sup>	Avg. Yield <sup>§</sup> (bu/ac)	Moisture at Harvest (%)	Test Weight (lbs/bu)
Integra 6915 TRE*	RR	TRE	266 A	12.6 A	60 B-D
Progeny PGY 2419TRE	RR	TRE	260 A	12.3 A	59 D
Dekalb DKC 119-30	RR	VT4P	255 A	12.9 A	60 BC
Dyna-Gro D60TC45RIB	RR	TRE	255 A	12.5 A	59 D
Pioneer P18216PWE	RR, LL, ENL, FOP	VT2P, HX1	254 A	12.7 A	60 B
Crow's 5859 TRE	RR	TRE	253 A	12.4 A	59 CD
Innvictis A1993T*	RR	TRE	251 A	12.3 A	60 B-D
Revere 1839 TC**	RR	TRE	249 A	12.1 A	60 B-D
Pioneer P17677YHR	RR, LL	YGCB, HX1	248 A	12.0 A	61 B
Dekalb DKC 68-35**	RR	VT2P	239 A	13.1 A	60 B-D
Innvictis A1792T	RR	TRE	237 A	12.4 A	62 A
<b>Trial Average</b>			252	12.5	60
<b>Trial Standard Error</b>			8	0.4	0
<b>Trial L.S.D.<sub>.05</sub></b>			N.S.	N.S.	1
<b>Trial C.V.</b>			4	4	1

† Hybrids that have any MS letter in common are not significantly different at the 5% level of probability.

\* Asterisks after a hybrid name indicate the number of preceding consecutive years in the top-performing "A" group.

‡ For a full description of abbreviated biotech traits, see table 4.

§ All yields are adjusted to 15.5% moisture.

|| Protein, Oil, and Starch on a dry weight basis.

Values highlighted in orange are above average, values highted in dark orange are in the upper 25%. MS letters highlighted in dark orange are in the "A group", indicating no statistical difference from the top-performing variety, for a given trait.